

## Control methods of *Botrytis cinerea*

Sergiu Fendrihan, Marian Lixandru, Sorina Dinu

Laboratory of Useful Organisms, Research Development Institute for Plant Protection,

Ion Ionescu de la Brad no 8, s. 1, Bucharest, ROMANIA

[ecologos23@yahoo.com](mailto:ecologos23@yahoo.com)

### ABSTRACT:

*Botrytis cinerea*, the agent of gray mold, are attacking more of 200 plant species and produce losses in postharvest products. Methods of control of this damaging fungus are reviewed in this article. Copper derived products are used and organic fungicide of few kind (boscalid, bezimidazole, carbendazim, propiconazole, and so on) as spray, some inorganic pesticide ZnO as nanoparticles. Organic natural derived substances produced by fungi *Penicillium brasilianum* *Trichoderma*, and bacteria *Paenibacillus* sp., can be use against *Botrytis*. The most interesting idea is the use of antagonists (bacteria and fungi) in biological control.

**Key words:** grey mold, chemical methods, copper sulphate, fungicides, biological control methods.

### INTRODUCTION

The gray mold is produced by *Botrytis cinerea*, attacking over 200 species of plants [1]. The authors find that an entire array of proteins are involved in virulence and their release depends on the pH of environment which means there are a permanent adjustment of their secretome, and the production are regulated at the transcriptional level. Other consider that the number of plants species attacked is about 500 [2] and produces economic losses of about 10 to 100 billion dollars worldwide. The species is characterized by a very diverse structure, with strains dependent of host, on geography, having some genetic clusters [3]. Generally speaking, the populations of *Botrytis* are characterized by differences in morphology, reproduction, virulence [4]. They are anyway are non-homogenous species having two groups I and II, the first lacking transposable elements Boty and Flipper [5]. Analyzing the data, it is a weak association between population structure and geographical area, and a considerable difference according to the origin of the host

plant [6]. Some strains look like to be more virulent than the others, and infection incidence is dependent of temperature, at 20°C is highest, and is lower at 30°C and 5°C, the data for mycelia growth and conidial germination are similar [7]. The researches reveal the heterogeneity of field population of *B. cinerea* as regarding morphology and reproduction, virulence and secondary metabolite production [4]. By molecular biological studies, some scientists revealed that *B. cinerea* produced changes in proteome, the accumulation of some protein, the thickening of cell wall production of lytic enzymes, can be mechanism of host defense against attack [8]. There is a different sensibility to different fungicides, depending on fungicide type and on strain of *B. cinerea*; the experiments performed in vitro showed a lower resistance to benzimidazole and carboximides and higher resistance to anilino pyrimidines [9]. Variations in population of the fungi results after point mutation changing protein structure related with the resistance [10]. The strains resistant to chemicals are the results of point

mutations and some multiple resistant strains which mechanism consists in mutations producing over expression of transporters systems members like ABC system. Another species is *B. fragariae*, attacking strawberry cultures [11] have local importance as phyto pathogen, and its presents a resistance to antifungal too. One of the elements of host (vineyard) which confers resistance to this pathogen is the wax layer of the grapes cuticula [12].

## METHODS OF CONTROL OF GREY MOLD

We review the methods for *B cinerea* control , in order to compare and to discuss the necessary experiments for further development.

### 2.1. Copper based products

Copper based products are used for hundreds of years in Plant protection. They are based on copper sulphate, copper carbonate, copper oxychloride, copper oxychloride sulfate, copper zinc chromate, copper dihydrazine sulphate, cuprous oxide, copper hydroxide [13].

### 2.2. Organic pesticides

A problem of applying pesticide is the resistance to fungicides of *Botrytis cinerea*. Experiments performed in the period 2008-2011 in Italy, showed resistance of isolate in several degrees, to aniline-pyrimidines, to boscalid and others including multiple fungicide resistance, due to incorrect use of them [14]. Benzimidazole azo phenol derivatives are efficient antifungals, like compound V-5 which shown to be effective against *F. graminearum*, *A. solani*, *V. mali*, *B. cinerea*, and *C. lunata* [15]. Currently against *Botrytis* are used fungicides like boscalid,

carbendazim, iprodione, pyrethanil and propiconazole and experiments showed the difference in their activity in inhibition of some metabolic processes [16]. ZnO, formulated as nanoparticles (NPs) can inhibits growth of *Botrytis cinerea* by affecting their functions and deformation of hyphae, at over 3mmol/L<sup>-1</sup> to *B. cinerea* [17]. For controlling of gray mold strains with multiple fungicides resistance in tomato green houses, and were used pesticides (fenxetamid and Signum- boscalid and pyraclostrobin) for control [18].

### 2.3. Bioactive substances

Generally we speak about natural origin compound with regulatory function, elaborated by different organisms in their antagonistic fight. *Penicillium brasilianum* produce isoroquefortine C, griseofulvin, ergosterol peroxide, 3 $\beta$ -hydroxy-(22E,24R)-ergosta-5,8,22-trien-7-one, cerevisterol and (22E,24R)-6 $\beta$ -methoxyergosta-7,22-diene-3 $\beta$ ,5 $\alpha$ -diol with bioactivity against fungi (*B. cinerea*, *Alternaria solani*, *Fusarium solani*, and so on) and human pathogenic bacteria [19]. Trichothecenes are produced by some fungi like *Trichoderma harzianum* CECT 2413 which intermediate compounds Trichodiene (TD) induce systemic defence [20]. Protocatehuic acid was isolated from *Paenibacillus elgii* HOA 73, and showed a potent antifungal effect against *Rhizoctonia solani* and *Botrytis cinerea* at MIC of 64  $\mu$ g ml<sup>-1</sup> [21]. Fatty acids foods coat are used in food industry, and were tested for their action against *B. cinerea*, the results showed the reduction of its germination and attack on grapes over 54-96% compared with the control lot. In the same time FC stimulates the yeasts present on grapes [22].

## 2.4. Biological controls

For biological control of this necrotrophic pathogen, were used different bacteria from *Bacillus* genus. Specialists from Algeria [23] isolated from extreme environments strains related with *B. amyloliquefaciens*, *B. atrophaeus* and *B. mohavensis* producing proteases and cellulases, and some chitinases acting against fungi and with supplementary growth stimulating activity. The same, for the strains of *Bacillus subtilis* B27 and B 29 which reduced attack to grapevine up to 50 -60%, due to chitinolytic enzymes [24]. *Bacillus amyloliquefaciens* subsp. *plantarum* strains IMV B-7404 and BIM B-439D demonstrated an antagonism against vascular bacteriosis agent and produced inhibition of *B. cinerea* and *Penicillium expansum* [25]. The *Bacillus* strains are able to produced lipopeptides which can be a tool for biocontrol of postharvest phytopathogens [26]. Yeasts isolated from grapes and wine, can be use as antagonists of *B. cinerea* and from about 591 isolates *Aureobasidium pullulans*, *Metschnikowia pulcherrima* and *Pichia guilliermondii* showed a good antagonistic effect even better than the *Candida oleophila* which is used usually for biocontrol [27]. Yeast *Candida sake* was effective against grey mould of tomato [18]. Other yeast strains with bio-control capacities are *Saccharomyces cerevisiae*, *Wickerhamomyces anomalus*, *Metschnikowia pulcherrima* *Aureobasidium pullulans*, strains of the first and the second, having the best antifungal activity against *Botrytis cinerea* [28]. *Trichoderma* spp. [29] has potential as biocontrol agent for *Botrytis cinerea*.

Some bacteria can protect plants from *B. cinerea* attack which synthesize anti-fungal metabolites, induce host resistance and are

competitive for nutrients with fungi [30]. A strain of *Brevibacillus brevis*, W4 [31], has a clear inhibitory effect on *B. cinerea*.

Actinomycetes are known as releasing precious secondary metabolites. *Streptomycin patents* LMM15 (its fermented substrate) can inhibits growth of grey mould and reduce lesions formed on tomato leaves and fruits by increasing content of proline and malon dialdehyde [32], providing the defense of plants.

Some fungi can defend the plants against grey mold too. *Trichoderma* sp and *Clonostachys* sp strains reduced mortality of mini cutting of *Eucalyptus globulus* in forest nurseries reducing disease in the same results like commercial fungicides [33]. Following experiments performed in vitro and in vivo, *Burkholderia cepacia* Cs5 demonstrated antagonistic capacity on fungus *B. cinerea* [34], being a possible future bio-fungicides.

## CONCLUSIONS

The fight against this necrotrophic fungi is a real challenge. The chemical methods of control are generally applied but it's affecting the environment despite all preventive measure taken and possible the consumer's safety. Practically a better method is the biological control, using antagonistic microorganisms. The examples e described in the present work, prove that this is really possible, being cheap, safe for man and environment and effective against *Botrytis cinerea* and possible other phytopathogenic fungi. There are still many aspects to be clarified by further researches.

**ACKNOWLEDGMENT**

We like to express sincere appreciation and deep gratitude to all participants in this work.

**REFERENCES**

- [1] Li B, Wang W, Zong Y, Qin G, Tian S. (2012) Exploring pathogenic mechanisms of *Botrytis cinerea* secretome under different ambient pH based on comparative proteomic analysis. *J Proteome Res*;11(8):4249-60.
- [2] Hua L, Yong C., Zhanquan Z., Boqiang L., Guozheng Q., Shiping T. (2018) Pathogenic mechanisms and control strategies of *Botrytis cinerea* causing post-harvest decay in fruits and vegetables, *Food Quality and Safety*, fyy016, <https://doi.org/10.1093/fqsafe/fyy016>.
- [3] Walker A.S. (2013) Diversité et adaptation aux fongicides des populations de *Botrytis cinerea*, agent de la pourriture grise. Sciences agricoles. These de doctorat, Université Paris Sud - Paris XI, 2013. Français
- [4] Schumacher J, Simon A, Cohrs KC, Traeger S, Porquier A, Dalmais B, Viaud M, Tudzynski B. (2015) The VELVET complex in the gray mold fungus *Botrytis cinerea*: impact of BcLAE1 on differentiation, secondary metabolism and virulence. *Mol Plant Microbe Interact*. 2015 Jan 27. [Epub ahead of print]
- [5] Fekete E., Fekete E., Irinyi L., Karaffa L., Árnýasi M., Asadollahi M, Sándor E. (2012) Genetic diversity of a *Botrytis cinerea* cryptic species complex in Hungary *Microbiological Research* 167:283– 291.
- [6] Walker AS, Gladieux P, Decognet V, Fermaud M, Confais J, Roudet J, Bardin M, Bout A, C Nicot P, Poncet C, Fournier E. (2015) Population structure and temporal maintenance of the multihost fungal pathogen *Botrytis cinerea*: causes and implications for disease management. *Environ Microbiol*;17(4):1261-74.
- [7] Ciliberti N, Fermaud M, Languasco L, Rossi V. (2015) Influence of Fungal Strain, Temperature, and Wetness Duration on Infection of Grapevine Inflorescences and Young Berry Clusters by *Botrytis cinerea*. *Phytopathology*;105(3):325-33.
- [8] Dadakova K, Havelkova M, Kurkova B, Tloková , Kasparovsky T, Zdrahal Z, Lochman J. (2015) Proteome and transcript analysis of *Vitis vinifera* cell cultures subjected to *Botrytis cinerea* infection. *J Proteomics*;119:143-53.
- [9] Bertetti D, Garibaldi A (2008) Resistance of *Botrytis cinerea* to fungicides in Italian vineyards. *Communications in agricultural and applied biological sciences*. 73. 273-82.
- [10] Chatzidimopoulos M, Psomopoulos F, Malandrakis EE, Ganopoulos I, Madesis P, Vellios EK and Drogoudi P (2016) Comparative Genomics of *Botrytis cinerea* Strains with Differential Multi-Drug Resistance. *Front. Plant Sci*. 7:554. doi: 10.3389/fpls.2016.00554.
- [11] Rupp S., Plesken C., Rumsey S., Dowling M., Schnabel G., Weber R.W. S., Hahn M. (2017) *Botrytis fragariae*, a New Species Causing Gray Mold on Strawberries, Shows High Frequencies of Specific and Efflux-Based Fungicide Resistance. *Appl Environ Microbiol*; 83(9): e00269-17
- [12] Herzog K, Rolf Wind and Reinhard Töpfer (2015) Impedance of the Grape Berry Cuticle as a Novel Phenotypic Trait to Estimate Resistance to *Botrytis Cinerea* Sensors; 15:12498-12512.
- [13] Van Zwieten M., Gordon Stovold, Lukas Van Zwieten (2007) Alternatives to Copper for Disease Control in the Australian Organic Industry A report for the Rural Industries Research and Development Corporation July 2007 RIRDC Publication No 07/110 RIRDC Project No DAN-208A
- [14] De Miccolis Angelini M.R., Rotolo C., Masiello M., Gerin D., Pollastro S., Faretra F. (2014) Occurrence of fungicide resistance in populations of *Botryotinia fuckeliana* (*Botrytis cinerea*) on table grape and strawberry in southern Italy *Pest Mag. Sci*;70 (12):1785-96.
- [15] Ke Y, Zhi X, Yu X, Ding G, Yang C, Xu H. (2014) Combinatorial synthesis of benzimidazole-azo-phenol derivatives as antifungal



- agents. Comb Chem High Throughput Screen.; 17(1):89-95.
- [16] Wang et al (2016) Metabolic activities of five botryticides against *Botrytis cinerea* examined using the Biolog FF MicroPlate. Sci Rep. 2016; 6: 31025. Published online 2016 Aug 5. doi: 10.1038/srep31025
- [17] He L, Yang Liu, Azlin Mustapha, Mengshi Lin (2011) Antifungal activity of zinc oxide nanoparticles against *Botrytis cinerea* and *Penicillium expansum* Microbiological Research 166 (2011) 207–215.
- [18] Rodríguez A, Acosta A, Rodríguez C. (2014) Fungicide resistance of *Botrytis cinerea* in tomato greenhouses in the Canary Islands and effectiveness of non-chemical treatments against gray mold. World J Microbiol Biotechnol.; 30(9):2397-406.
- [19] Tang HY, Zhang Q, Li H, Gao JM. (2015) Antimicrobial and allelopathic metabolites produced by *Penicillium brasilianum*. Nat Prod Res.; 29(4):345-8.
- [20] Malmierca MG, McCormick SP, Cardoza RE, Alexander NJ, Monte E, Gutiérrez S. (2014) Production of trichodiene by *Trichoderma harzianum* alters the perception of this biocontrol strain by plants and antagonized fungi. Environ Microbiol. 2014 May 12. doi: 10.1111/1462-2920.12506. [Epub ahead of print]
- [21] Nguyen XH, Naing KW, Lee YS, Moon JH, Lee JH, Kim KY. (2015) Isolation and characteristics of protocatechuic acid from *Paenibacillus elgii* HOA73 against *Botrytis cinerea* on strawberry fruits. J Basic Microbiol.; 55(5):625-34.
- [22] Calvo-Garrido C, Elmer PA, Parry FJ, Viñas I, Usall J, Torres R, Agnew RH, Teixidó N. (2014) Mode of action of a fatty acid-based natural product to control *Botrytis cinerea* in grapes. J Appl Microbiol.; 116(4):967-79.
- [23] Ait-Kaki A, Kacem-Chaouche N, Ongena M, Kara-Ali M, Dehimat L, Kahlat K, Thonart P. (2014) In vitro and in vivo characterization of plant growth promoting *Bacillus* strains isolated from extreme environments of Eastern Algeria. Appl Biochem Biotechnol.; 172(4):1735-46.
- [24] Maachia B, Rafik E, Chérif M, Nandal P, Mohapatra T, Bernard P. (2015) Biological control of the grapevine diseases 'grey mold' and 'powdery mildew' by *Bacillus* B27 and B29 strains. Indian J Exp Biol.; 53(2):109-15.
- [25] Avdeeva LV, Dragovoz IV, Korzh IuV, Leonova NO, Iutinskaia GA, Berezhnaia AV, Kuptsov VN, Mandrik MN, Kolomiets ÉI (2014) [Antagonistic activity of *Bacillus amyloliquefaciens* subsp. *plantarum* IMV B-7404 and BIM B-439D strains towards pathogenic bacteria and micromycetes] [Article in Russian] Mikrobiol Z.; 76(6):27-33.
- [26] Pretorius D., van Rooyen J., Clarke K.G. (2015) Enhanced production of antifungal lipopeptides by *Bacillus amyloliquefaciens* for biocontrol of postharvest disease. N Biotechnol.; 32(2):243-52
- [27] Raspor P., Mikli-Milek D., Avbelj M., Cadez N. (2010) Biocontrol of Grey Mould Disease on Grape Caused by *Botrytis cinerea* with Autochthonous Wine Yeasts. Food Technol. Biotechnol. 48 (3) 336–343.
- [28] Parafati L, Vitale A, Restuccia C, Cirvilleri G. (2015) Biocontrol ability and action mechanism of food-isolated yeast strains against *Botrytis cinerea* causing post-harvest bunch rot of table grape. Food Microbiol.; 47:85-92.
- [29] Vos CM, De Cremer K, Cammue BP, De Coninck B (2015) The toolbox of *Trichoderma* spp. in the biocontrol of *Botrytis cinerea* disease. Mol Plant Pathol.; 16(4):400-12. doi: 10.1111/mpp.12189. Epub 2014 Oct 22.
- [30] Haidar R., Fermaud M, Calvo-Garrido C., Roudet J., Deschamps A. (2016) Modes of action for biological control of *Botrytis cinerea* by antagonistic bacteria. Phytopathologia Mediterranea (2016) 55, 3, 301–322.
- [31] Yang C, Zhang X, Shi G, Zhao H, Chen L, Tao K and Hou T (2011) Isolation and identification of endophytic bacterium W4 against tomato *Botrytis cinerea* and antagonistic activity stability. African Journal of Microbiology Research; 5(2): 131.
- [32] Lian Q., Zhang J., Gan L., Ma Q., Zong Z., Wang Y. (2017) The Biocontrol Efficacy of

*Streptomyces pratensis* LMM15 on *Botrytis cinerea* in Tomato. Hindawi BioMed Research International Volume 2017, Article ID 9486794, 11 pages  
<https://doi.org/10.1155/2017/9486794>

- [33] Zaldúa S., Sanfuentes E. (2010) Control of *Botrytis cinerea* in *Eucalyptus globulus* mini-cuttings using *Clonostachys* and *Trichoderma* strains. Chilean J. of Agric. Res. 70 (4):576-582.
- [34] Kilani-Feki O., Jaoua S. (2011) Biological control of *Botrytis cinerea* using the antagonistic and endophytic *Burkholderia cepacia* Cs5 for vine plantlet protection. Can. J. Microbiol. 57: 896–901.